

51798

MEASUREMENT OF POST-VOIDING RESIDUAL URINE AND/OR  
BLADDER CAPACITY BY ULTRASOUND, NON-IMAGING

ICD-10 Codes that Support Medical Necessity

Group 1 Paragraph: It is the responsibility of the provider to code to the highest level specified in the ICD-10-CM. The correct use of an ICD-10-CM code listed below does not assure coverage of a service. The service must be reasonable and necessary in the specific case and must meet the criteria specified in this determination.

Group 1 Codes:

<u>ICD-10 Code</u>	<u>Description</u>
N31.0 - N31.2	Uninhibited neuropathic bladder, not elsewhere classified - Flaccid neuropathic bladder, not elsewhere classified
N31.9	Neuromuscular dysfunction of bladder, unspecified
N32.0	Bladder-neck obstruction
N39.3	Stress incontinence (female) (male)
N39.41 - N39.46	Urge incontinence - Mixed incontinence
N39.490	Overflow incontinence
N39.498	Other specified urinary incontinence
N40.1	Enlarged prostate with lower urinary tract symptoms
R30.0	Dysuria
R30.1	Vesical tenesmus
R30.9	Painful micturition, unspecified
R32	Unspecified urinary incontinence
R33.0	Drug induced retention of urine
R33.8	Other retention of urine
R33.9	Retention of urine, unspecified
R34	Anuria and oliguria
R35.0	Frequency of micturition
R35.1	Nocturia
R35.8	Other polyuria
R36.0	Urethral discharge without blood
R36.9	Urethral discharge, unspecified
R39.0	Extravasation of urine
R39.11 - R39.16	Hesitancy of micturition - Straining to void
R39.19	Other difficulties with micturition
R39.2	Extrarenal uremia
R39.81	Functional urinary incontinence
R39.89	Other symptoms and signs involving the genitourinary system